


## On birecurrent for some tensors in various Finsler spaces

Alaa A. Abdallah<sup>a</sup>, Ahmed A. Hamoud<sup>b</sup>, A. Navlekar<sup>c</sup>, Kirtiwant Ghadle<sup>d</sup>,  
Basel Hardan<sup>e</sup>, Homan Emadifar<sup>f\*</sup> , Masoumeh Khademi<sup>f</sup>

<sup>a</sup>Department of Mathematics, Abyan University, Abyan, Yemen.

E-mail: maths.aab@bamu.ac.in

<sup>b</sup>Department of Mathematics, Taiz University, Taiz P.O. Box 6803, Yemen

E-mail: ahmed.hamoud@taiz.edu.ye

<sup>c</sup>Department of Mathematics, Pratishthan Mahavidyalaya, Paithan, India.

E-mail: dr.navlekar@gmail.com

<sup>d</sup>Department of Mathematics, Dr. Babasaheb Ambedkar Marathwada

University, Aurangabad, India.

E-mail: ghadle.maths@bamu.ac.in

<sup>e</sup>Department of Mathematics, Abyan University, Abyan, Yemen.

E-mail: bassil2003@gmail.com

<sup>f</sup>Department of Mathematics, Hamedan Branch, Islamic Azad University,

Hamedan, Iran.

E-mail: homan\_emadi@yahoo.com(H.E), dr.amonaft@gmail.com(M.KH)

**Abstract.** The  $\mathfrak{BC}$ - recurrent Finsler space introduced by Alaa et al. [1]. Now in this paper, we introduce and extend  $\mathfrak{BC}$ - birecurrent Finsler space by using some properties of different spaces. We study the relationship between Cartan's second curvature tensor  $P_{jkh}^i$  and  $(h)hv$ - torsion tensor  $C_{jk}^i$  in sense of Berwald. Additionally, the necessary and sufficient condition for some tensors which satisfy birecurrence property will be discuss in different spaces. Four theorems have been established and proved.

**Keywords:**  $\mathfrak{BC}$ - birecurrent space, birecurrence property,  $P2$ -like space,

---

\*Corresponding Author

AMS 2020 Mathematics Subject Classification: 53B40, 53C42, 53C60

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

Copyright © 2023 The Author(s). Published by University of Mohaghegh Ardabili

$P^*$ -space, generalized  $P$ -reducible space.

## 1. Introduction

The tensors which satisfy a birecurrence property in Finsler spaces has been discussed by the Finslerian geometers. The concept of  $C$ -birecurrent space in sense of Cartan and Berwald were studied by Pandey and Verma [20] and Sarangi and Goswami [13], respectively. Saleem [6] discussed  $C^h$ -generalized birecurrent space and  $C^h$ -special generalized birecurrent space. Pandey and Verma [20], Otman [9], Hanballa [8], Alqufail et al. [14] and Dikshit [23] introduced  $C^h$ -birecurrent space,  $\mathfrak{B}P$ -birecurrent space,  $\mathfrak{B}K$ -birecurrent space,  $K^h$ -birecurrent space and  $R^h$ -birecurrent space, respectively. Also, Qasem and Hanballa [10] studied  $K^h$ -generalized birecurrent space.

In the same vein, Saleem and Abdallah [7] introduced the  $U^h$ -birecurrent Finsler space and discussed the necessary and sufficient condition for some tensors which satisfy the birecurrence property.

Regarding to special spaces of Finsler space, Pandey and Dikshit [21] discussed  $P^*$ - and  $P$ -reducible Finsler space of recurrent curvature tensor, Otman [9] studied the properties of  $P2$ -like space and  $P^*$ -space in  $P^h$ -birecurrent space. In addition, Saleem [6] studied  $P2$ -like-generalized birecurrent space and  $P2$ -like- $C^h$ -special generalized birecurrent. Further, Saxena and Swarup [22] used  $P$ -reducibility condition in spacial Finsler spaces. Recently, the properties of  $P2$ -like space,  $P^*$ -space and generalized  $P$ -reducible space in generalized  $\mathfrak{B}P$ -recurrent space have been studied by [2, 3]. The main idea of this paper to concentrate on obtaining the necessary and sufficient condition for  $P_{jkh}^i, P_{ijkh}, P_{kh}^i, P_{jk}, P_k$  and  $P$  which satisfy birecurrence property in various spaces.

## 2. Preliminaries

In this section, important concept of Finsler geometry will be given in this paper. An  $n$ -dimensional space  $X_n$  equipped with a function  $F(x, y)$  that denoted by  $F_n = (X_n, F(x, y))$  called a Finsler space if the function  $F(x, y)$  satisfying the request conditions [5, 12, 15, 24].

Matsumoto [18] introduced the  $(h)hv$ -torsion tensor  $C_{ijk}$  that is positively homogeneous of degree  $-1$  in  $y^j$  and defined by

$$C_{ijk} = \frac{1}{2} \dot{\partial}_i g_{jk} = \frac{1}{4} \dot{\partial}_i \dot{\partial}_j \dot{\partial}_k F^2.$$

By using Euler's theorem on homogeneous function, we get

$$a) C_{ijk} y^i = C_{kij} y^i = C_{jki} y^i = 0 \text{ and } b) C_{jk}^i y^j = C_{kj}^i y^j = 0, \quad (2.1)$$

where  $C_{jk}^i$  is called associate tensor of the tensor  $C_{ijk}$ , these tensors are defined by

$$a) C_{ik}^h = C_{ijk}g^{hj}, \quad b) C_{ji}^i = C_j \quad \text{and} \quad c) C_k y^k = C, \quad (2.2)$$

The unit vector  $l^i$  and the associative vector  $l_i$  with the direction of  $y^i$  are given by

$$a) l^i = \frac{y^i}{F} \quad \text{and} \quad b) l_i = \frac{y_i}{F}. \quad (2.3)$$

Berwald covariant derivative  $\mathfrak{B}_k T_j^i$  of an arbitrary tensor field  $T_j^i$  with respect to  $x^k$  is given by [12]

$$\mathfrak{B}_k T_j^i = \partial_k T_j^i - (\dot{\partial}_r T_j^i) G_k^r + T_j^r G_{rk}^i - T_r^i G_{jk}^r.$$

Berwald covariant derivative of the vector  $y^i$  vanish identically, i.e.

$$\mathfrak{B}_k y^i = 0. \quad (2.4)$$

The tensor  $P_{jkh}^i$  is called  $hv$ -curvature tensor (Cartan's second curvature tensor) is positively homogeneous of degree -1 in  $y^i$  and defined by [12]

$$P_{jkh}^i = C_{kh|j}^i - g^{ir} C_{jkh|r} + C_{jk}^r P_{rh}^i - P_{jh}^r C_{rk}^i, \quad (2.5)$$

which satisfies the relation

$$P_{jkh}^i y^j = \Gamma_{jkh}^{*i} y^j = P_{kh}^i = C_{kh|r}^i y^r, \quad (2.6)$$

where  $P_{kh}^i$  is  $(v)hv$ -torsion tensor which satisfies

$$P_{kh}^i = P_{rkh} g^{ir}, \quad (2.7)$$

where  $P_{rkh}$  is called associative tensor for  $v(hv)$ -torsion tensor.

$P$ -Ricci tensor  $P_{jk}$ , curvature vector  $P_k$  and curvature scalar  $P$  of Cartan's second curvature tensor are given by

$$a) P_{jk} = P_{jki}^i, \quad b) P_{ki}^i = P_k \quad \text{and} \quad c) P = P_k y^k, \quad (2.8)$$

respectively.

**Definition 2.1.** A Finsler space  $F_n$  is called a  $P2$ -like space if the Cartan's second curvature tensor  $P_{jkh}^i$  is characterized by the condition [18]

$$P_{jkh}^i = \varphi_j C_{kh}^i - \varphi^i C_{jkh}, \quad (2.9)$$

where  $\varphi_j$  and  $\varphi^i$  are non-zero covariant and contravariant vectors field, respectively.

**Definition 2.2.** A Finsler space  $F_n$  is called a  $P^*$ -Finsler space if the  $(v)hv$ -torsion tensor  $P_{kh}^i$  is characterized by the condition [11]

$$P_{kh}^i = \varphi C_{kh}^i, \quad (2.10)$$

where  $P_{jkh}^i y^j = P_{kh}^i = C_{kh|s}^i y^s$ .

**Definition 2.3.** A Finsler space  $F_n$  is called a generalized  $P$ -reducible space if the associate tensor  $P_{jkh}$  of  $(v)hv$ -torsion tensor  $P_{kh}^i$  is characterized by the condition [19, 25]

$$P_{jkh} = \lambda C_{jkh} + \vartheta(h_{jk}C_h + h_{kh}C_j + h_{hj}C_k), \quad (2.11)$$

where  $\lambda$  and  $\vartheta$  are scalar vectors positively homogeneous of degree one in  $y^j$  and  $h_{jk}$  is the angular metric tensor.

**Definition 2.4.** Let the current coordinates in the tangent space at the point  $x_0$  be  $x^i$ , then the indicatrix  $I_{n-1}$  is a hypersurface defined by [12]  $F(x_0, x^i) = 1$  or by the parametric form defined by  $x^i = x^i(u^a)$ ,  $a = 1, 2, \dots, n-1$ .

**Definition 2.5.** The projection of any tensor  $T_j^i$  on indicatrix  $I_{n-1}$  given by [12, 16]

$$p.T_j^i = T_b^a h_a^i h_j^b, \quad (2.12)$$

where

$$h_c^i = \delta_c^i - l^i l_c. \quad (2.13)$$

The projection of the vector  $y^i$ , the unit vector  $l^i$  and the metric tensor  $g_{ij}$  on the indicatrix are given by  $p.y^i = 0$ ,  $p.l^i = 0$  and  $p.g_{ij} = h_{ij}$ , where  $h_{ij} = g_{ij} - l_i l_j$ .

### 3. On $\mathfrak{BC}$ -Birecurrent Space

In this section, we find the condition for different tensors which behave as birecurrent in  $\mathfrak{BC}$ -birecurrent space. Matsumoto [17] introduced a Finsler space which the  $(h)hv$ -torsion tensor  $C_{ijk}$  and its associate tensor  $C_{jk}^i$  satisfy the recurrence property in sense of Cartan. This space is called  $C^h$ -recurrent space and characterized by the conditions

$$a) C_{kh|m}^i = \lambda_m C_{kh}^i \text{ and } b) C_{jkh|m} = \lambda_m C_{jkh}. \quad (3.1)$$

Alaa et al. [1] introduced  $\mathfrak{BC} - RF_n$  which is characterized by the conditions

$$a) \mathfrak{B}_m C_{kh}^i = \lambda_m C_{kh}^i \text{ and } b) \mathfrak{B}_m C_{jkh} = \lambda_m C_{jkh}. \quad (3.2)$$

Sarangi and Goswami [13] introduced a Finsler space which the  $(h)hv$ -torsion tensor  $C_{ijk}$  and its associate tensor  $C_{jk}^i$  satisfy the birecurrence property in sense of Berwald and called it  $C$ -birecurrent space. Let us denote to this space briefly by a  $\mathfrak{BC} - BRF_n$ . This space characterized by the conditions

$$a) \mathfrak{B}_l \mathfrak{B}_m C_{kh}^i = a_{lm} C_{kh}^i \text{ and } b) \mathfrak{B}_l \mathfrak{B}_m C_{jkh} = a_{lm} C_{jkh}, \quad (3.3)$$

where  $a_{lm} = \mathfrak{B}_l \lambda_m + \lambda_l \lambda_m$ . Using eq. (3.1) in (2.5), we get

$$P_{jkh}^i = \lambda_j C_{kh}^i - \lambda^i C_{jkh} + C_{jk}^r P_{rh}^i - C_{rk}^i P_{jh}^r, \quad (3.4)$$

where  $\lambda^i = \lambda_r g^{ir}$ .

In next theorem we get the necessary and sufficient condition for some tensors which behave as birecurrent tensor in  $\mathfrak{BC} - BRF_n$ .

**Theorem 3.1.** *In  $\mathfrak{BC} - BRF_n$ , Cartan's second curvature tensor  $P_{jkh}^i$ , torsion tensor  $P_{kh}^i$ ,  $P$ -Ricci tensor  $P_{jk}$ , curvature vector  $P_k$  and curvature scalar  $P$  satisfy the birecurrence property if and only if*

$$\begin{aligned} & \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_{kh}^i \\ & - \{(\mathfrak{B}_l \mathfrak{B}_m \lambda^i) - (\mathfrak{B}_m \lambda^i) \lambda_l - (\mathfrak{B}_l \lambda^i) \lambda_m\} C_{jkh} \\ & + \{\lambda_m (\mathfrak{B}_l P_{rh}^i) + \lambda_l (\mathfrak{B}_m P_{rh}^i) + (\mathfrak{B}_l \mathfrak{B}_m P_{rh}^i)\} C_{jk}^r \\ & - \{\lambda_m (\mathfrak{B}_l P_{jh}^r) + \lambda_l (\mathfrak{B}_m P_{jh}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{jh}^r)\} C_{rk}^i = 0, \end{aligned} \quad (3.5)$$

$$\begin{aligned} & \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_{kh}^i y^j \\ & - \{\lambda_m (\mathfrak{B}_l P_{jh}^r) + \lambda_l (\mathfrak{B}_m P_{jh}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{jh}^r)\} C_{rk}^i y^j = 0, \end{aligned} \quad (3.6)$$

$$\begin{aligned} & \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_k \\ & - \{(\mathfrak{B}_l \mathfrak{B}_m \lambda^i) - (\mathfrak{B}_m \lambda^i) \lambda_l - (\mathfrak{B}_l \lambda^i) \lambda_m\} C_{jki} \\ & + \{\lambda_m (\mathfrak{B}_l P_r) + \lambda_l (\mathfrak{B}_m P_r) + (\mathfrak{B}_l \mathfrak{B}_m P_r)\} C_{jk}^r \\ & - \{\lambda_m (\mathfrak{B}_l P_{ji}^r) + \lambda_l (\mathfrak{B}_m P_{ji}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{ji}^r)\} C_{rk}^i = 0, \end{aligned} \quad (3.7)$$

$$\begin{aligned} & \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_k y^j \\ & - \{\lambda_m (\mathfrak{B}_l P_{ji}^r) + \lambda_l (\mathfrak{B}_m P_{ji}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{ji}^r)\} C_{rk}^i y^j = 0, \end{aligned} \quad (3.8)$$

and

$$\{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C y^j = 0, \quad (3.9)$$

respectively.

Proof. Taking  $\mathfrak{B}$ -covariant derivative for eq. (3.4) twice with respect to  $x^m$  and  $x^l$ , respectively, using eqs. (3.2) and (3.3) in the resulting equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jkh}^i &= a_{lm} (\lambda_j C_{kh}^i - \lambda^i C_{jkh} + C_{jk}^r P_{rh}^i - C_{rk}^i P_{jh}^r) \\ &+ \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_{kh}^i \\ &- \{(\mathfrak{B}_l \mathfrak{B}_m \lambda^i) - (\mathfrak{B}_m \lambda^i) \lambda_l - (\mathfrak{B}_l \lambda^i) \lambda_m\} C_{jkh} \\ &+ \{\lambda_m (\mathfrak{B}_l P_{rh}^i) + \lambda_l (\mathfrak{B}_m P_{rh}^i) + (\mathfrak{B}_l \mathfrak{B}_m P_{rh}^i)\} C_{jk}^r \\ &- \{\lambda_m (\mathfrak{B}_l P_{jh}^r) + \lambda_l (\mathfrak{B}_m P_{jh}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{jh}^r)\} C_{rk}^i. \end{aligned}$$

Using eq. (3.4) in above equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jkh}^i &= a_{lm} P_{jkh}^i + \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_{kh}^i \\ &- \{(\mathfrak{B}_l \mathfrak{B}_m \lambda^i) - (\mathfrak{B}_m \lambda^i) \lambda_l - (\mathfrak{B}_l \lambda^i) \lambda_m\} C_{jkh} \\ &+ \{\lambda_m (\mathfrak{B}_l P_{rh}^i) + \lambda_l (\mathfrak{B}_m P_{rh}^i) + (\mathfrak{B}_l \mathfrak{B}_m P_{rh}^i)\} C_{jk}^r \\ &- \{\lambda_m (\mathfrak{B}_l P_{jh}^r) + \lambda_l (\mathfrak{B}_m P_{jh}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{jh}^r)\} C_{rk}^i. \end{aligned} \quad (3.10)$$

This shows that

$$\mathfrak{B}_l \mathfrak{B}_m P_{jkh}^i = a_{lm} P_{jkh}^i. \quad (3.11)$$

if and only if eq. (3.5) holds.

Transvecting eq. (3.10) by  $y^j$ , using (2.1), (2.4) and (2.6), we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{kh}^i &= a_{lm} P_{kh}^i + \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_{kh}^i y^j \\ &\quad - \{\lambda_m (\mathfrak{B}_l P_{jh}^r) + \lambda_l (\mathfrak{B}_m P_{jh}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{jh}^r)\} C_{rk}^i y^j \end{aligned} \quad (3.12)$$

This shows that

$$\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i. \quad (3.13)$$

if and only if eq. (3.6) holds.

Contracting the indices  $i$  and  $h$  in eq. (3.10), using (2.2) and (2.8), we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jk} &= a_{lm} P_{jk} + \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_k \\ &\quad - \{(\mathfrak{B}_l \mathfrak{B}_m \lambda^i) - (\mathfrak{B}_m \lambda^i) \lambda_l - (\mathfrak{B}_l \lambda^i) \lambda_m\} C_{jki} \\ &\quad + \{\lambda_m (\mathfrak{B}_l P_r) + \lambda_l (\mathfrak{B}_m P_r) + (\mathfrak{B}_l \mathfrak{B}_m P_r)\} C_{jk}^r \\ &\quad - \{\lambda_m (\mathfrak{B}_l P_{ji}^r) + \lambda_l (\mathfrak{B}_m P_{ji}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{ji}^r)\} C_{rk}^i. \end{aligned} \quad (3.14)$$

This shows that

$$\mathfrak{B}_l \mathfrak{B}_m P_{jk} = a_{lm} P_{jk}. \quad (3.15)$$

if and only if eq. (3.7) holds.

Contracting the indices  $i$  and  $h$  in eq. (3.12), using (2.2) and (2.8), we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_k &= a_{lm} P_k + \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C_k y^j \\ &\quad - \{\lambda_m (\mathfrak{B}_l P_{ji}^r) + \lambda_l (\mathfrak{B}_m P_{ji}^r) + (\mathfrak{B}_l \mathfrak{B}_m P_{ji}^r)\} C_{rk}^i y^j \end{aligned} \quad (3.16)$$

This shows that

$$\mathfrak{B}_l \mathfrak{B}_m P_k = a_{lm} P_k \quad (3.17)$$

if and only if eq. (3.8) holds.

Transvecting eq. (3.16) by  $y^k$ , using (2.2), (2.4) and (2.8), we get

$$\mathfrak{B}_l \mathfrak{B}_m P = a_{lm} P + \{(\mathfrak{B}_l \mathfrak{B}_m \lambda_j) + (\mathfrak{B}_m \lambda_j) \lambda_l + (\mathfrak{B}_l \lambda_j) \lambda_m\} C y^j \quad (3.18)$$

This shows that

$$\mathfrak{B}_l \mathfrak{B}_m P = a_{lm} P. \quad (3.19)$$

if and only if eq. (3.9) holds.

Consequently, from eqs. (3.11), (3.13), (3.15), (3.17) and (3.19), we deduce that the behavior of  $P_{jkh}^i$ ,  $P_{kh}^i$ ,  $P_{jk}$ ,  $P_k$  and  $P$  in  $\mathfrak{BC} - BRF_n$  as birecurrent if and only if eqs. (3.5), (3.6), (3.7), (3.8) and (3.9), respectively hold. Hence, we have proved this theorem.

#### 4. Special Spaces of $\mathfrak{BC}$ –Birecurrent Space

In this section, we merge the  $\mathfrak{BC}$ – birecurrent space with particular spaces of Finsler space to get new spaces.

##### 4.1. A $P2$ –Like $\mathfrak{BC}$ –Birecurrent Space.

**Definition 4.1.** *The  $\mathfrak{BC}$ –birecurrent space which is  $P2$ –like space, i.e. satisfies the condition (2.9), will be called a  $P2$ –like  $\mathfrak{BC}$ –birecurrent space and will be denoted briefly by  $P2$ –like– $\mathfrak{BC} - BRF_n$ .*

In next theorem we get the necessary and sufficient condition for some tensors which behave as birecurrent tensor in  $P2$ –like– $\mathfrak{BC} - BRF_n$ .

**Theorem 4.2.** *In  $P2$ –like– $\mathfrak{BC} - BRF_n$ , Cartan's second curvature tensor  $P_{jkh}^i$ , torsion tensor  $P_{kh}^i$ ,  $P$ –Ricci tensor  $P_{jk}$  and curvature vector  $P_k$  satisfy the birecurrence property if and only if*

$$\begin{aligned} & \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_{kh}^i \\ & - \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta^i) - (\mathfrak{B}_m \vartheta^i) \lambda_l - (\mathfrak{B}_l \vartheta^i) \lambda_m\} C_{jkh} = 0, \end{aligned} \quad (4.1)$$

$$\{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_{kh}^i y^j = 0, \quad (4.2)$$

$$\begin{aligned} & \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_k \\ & - \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta^i) - (\mathfrak{B}_m \vartheta^i) \lambda_l - (\mathfrak{B}_l \vartheta^i) \lambda_m\} C_{jki} = 0 \end{aligned} \quad (4.3)$$

and

$$\{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_k y^j = 0. \quad (4.4)$$

respectively.

Proof. Taking  $\mathfrak{B}$ – covariant derivative for the condition (2.9) twice with respect to  $x^m$  and  $x^l$ , respectively, using eqs. (3.2) and (3.3) in the resulting equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jkh}^i &= a_{lm} (\vartheta_j C_{kh}^i - \vartheta^i C_{jkh}) \\ &+ \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_{kh}^i \\ &- \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta^i) - (\mathfrak{B}_m \vartheta^i) \lambda_l - (\mathfrak{B}_l \vartheta^i) \lambda_m\} C_{jkh}. \end{aligned}$$

Using the condition (2.9) in above equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jkh}^i &= a_{lm} P_{jkh}^i + \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_{kh}^i \\ &- \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta^i) - (\mathfrak{B}_m \vartheta^i) \lambda_l - (\mathfrak{B}_l \vartheta^i) \lambda_m\} C_{jkh}. \end{aligned} \quad (4.5)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P_{jkh}^i = a_{lm} P_{jkh}^i$  if and only if eq. (4.1) holds.

Transvecting eq. (4.5) by  $y^j$  using (2.1), (2.4) and (2.6), we get

$$\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i + \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_{kh}^i y^j. \quad (4.6)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i$  if and only if eq. (4.2) holds.

Contracting the indices  $i$  and  $h$  in eq. (4.5), using (2.2) and (2.8), we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jk} &= a_{lm} P_{jk} + \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_k \\ &\quad - \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta^i) - (\mathfrak{B}_m \vartheta^i) \lambda_l - (\mathfrak{B}_l \vartheta^i) \lambda_m\} C_{jki}. \end{aligned} \quad (4.7)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P_{jk} = a_{lm} P_{jk}$  if and only if eq. (4.3) holds.

Contracting the indices  $i$  and  $h$  in eq. (4.6), using (2.2) and (2.8), we get

$$\mathfrak{B}_l \mathfrak{B}_m P_k = a_{lm} P_k + \{(\mathfrak{B}_l \mathfrak{B}_m \vartheta_j) + (\mathfrak{B}_m \vartheta_j) \lambda_l + (\mathfrak{B}_l \vartheta_j) \lambda_m\} C_k y^j \quad (4.8)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P_k = a_{lm} P_k$  if and only if eq. (4.4) holds.

Consequently, from previous equations we proved that the behavior of  $P_{jkh}^i$ ,  $P_{kh}^i$ ,  $P_{jk}$  and  $P_k$  in  $P2$ -like- $\mathfrak{BC} - BRF_n$  as birecurrent if and only if eqs. (4.1), (4.2), (4.3) and (4.4), respectively hold. Hence, we have proved this theorem.

#### 4.2. A $P^* - \mathfrak{BC}$ -Birecurrent Space.

**Definition 4.3.** *The  $\mathfrak{BC}$ -birecurrent space which is  $P^*$ -space, i.e. satisfies the condition (2.10), will be called a  $P^* - \mathfrak{BC}$ -birecurrent space and will be denoted briefly by  $P^* - \mathfrak{BC} - BRF_n$ .*

In next theorem we get the necessary and sufficient condition for some tensors which behave as recurrent tensor in  $P^* - \mathfrak{BC} - BRF_n$ .

**Theorem 4.4.** *In  $P^* - \mathfrak{BC} - BRF_n$ , the torsion tensor  $P_{kh}^i$ , curvature vector  $P_k$  and curvature scalar  $P$  satisfy the birecurrence property if and only if*

$$[\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C_{kh}^i = 0, \quad (4.9)$$

$$[\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C_k = 0 \quad (4.10)$$

and

$$[\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C = 0, \quad (4.11)$$

respectively.

Proof. Taking  $\mathfrak{B}$ -covariant derivative for the condition (2.10) twice with respect to  $x^m$  and  $x^l$ , respectively, using eqs.(3.2) and (3.3) in the resulting equation, we get

$$\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = \vartheta a_{lm} C_{kh}^i + [\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C_{kh}^i.$$

Using the condition (2.10) in above equation, we get

$$\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i + [\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C_{kh}^i \quad (4.12)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i$  if and only if eq. (4.9) holds.

Contracting the indices  $i$  and  $h$  in eq. (4.12), using (2.2) and (2.8), we get

$$\mathfrak{B}_l \mathfrak{B}_m P_k = a_{lm} P_k + [\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C_k \quad (4.13)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P_k = a_{lm} P_k$  if and only if eq. (4.10) holds.



Transvecting eq. (4.13) by  $y^k$ , using (2.2) and (2.8), we get

$$\mathfrak{B}_l \mathfrak{B}_m P = a_{lm} P + [\mathfrak{B}_l \mathfrak{B}_m \vartheta + \lambda_l (\mathfrak{B}_m \vartheta) + \lambda_m (\mathfrak{B}_l \vartheta)] C \quad (4.14)$$

This shows that  $\mathfrak{B}_l \mathfrak{B}_m P = a_{lm} P$  if and only if eq. (4.11) holds.

Consequently, from previous equations we proved that the behavior of  $P_{kh}^i$ ,  $P_k$  and  $P$  in  $P^* - \mathfrak{B}C - BRF_n$  as birecurrent if and only if eqs. (4.9), (4.10) and (4.11), respectively hold. Hence, we have proved this theorem.

#### 4.3. A $P$ -Reducible $-\mathfrak{B}C$ -Birecurrent Space.

**Definition 4.5.** *The  $\mathfrak{B}C$ -birecurrent space which is generalized  $P$ -reducible space, i.e. satisfies the condition (2.11), will be called a  $P$ -reducible  $-\mathfrak{B}C$ -birecurrent space and will be denoted briefly by  $P$ -reducible  $-\mathfrak{B}C - BRF_n$ .*

In next theorem we get the necessary and sufficient condition for some tensors which be non-vanishing in  $P$ -reducible  $-\mathfrak{B}C - BRF_n$ .

**Theorem 4.6.** *In  $P$ -reducible  $-\mathfrak{B}C - BRF_n$ , Berwald's covariant derivative of the second order for the tensors  $\vartheta(h_k^i C_h + h_{kh} C^i + h_h^i C_k)$  and  $\vartheta(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k)$  are given by*

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m [\vartheta(h_k^i C_h + h_{kh} C^j + h_h^i C_k)] &= a_{lm} \vartheta(h_k^i C_h + h_{kh} C^i + h_h^i C_k) \quad (4.15) \\ &- [\mathfrak{B}_l \mathfrak{B}_m \lambda + (\mathfrak{B}_m \lambda) \lambda_l + (\mathfrak{B}_l \lambda) \lambda_m] C_{kh}^i \end{aligned}$$

and

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m [\vartheta(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k)] &= a_{lm} \vartheta(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k) \\ &- [\mathfrak{B}_l \mathfrak{B}_m \lambda + (\mathfrak{B}_m \lambda) \lambda_l + (\mathfrak{B}_l \lambda) \lambda_m] C_{jkh} \end{aligned} \quad (4.16)$$

if and only if the torsion tensor  $P_{kh}^i$  and associate torsion tensor  $P_{jkh}$  satisfy the birecurrence property, respectively.

Proof. Transvecting the condition (2.11) by  $g^{ij}$ , using (2.7) and (2.2), we get

$$P_{kh}^i = \lambda C_{kh}^i + \vartheta(h_k^i C_h + h_{kh} C^i + h_h^i C_k), \quad (4.17)$$

where  $h_k^i = g^{ij} h_{jk}$  and  $C^i = g^{ij} C_j$ .

Taking  $\mathfrak{B}$ -covariant derivative for the condition (4.17) twice with respect to  $x^m$  and  $x^l$  respectively, using eqs. (3.2) and (3.3) in the resulting equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{kh}^i &= \lambda a_{lm} C_{kh}^i + [\mathfrak{B}_l \mathfrak{B}_m \lambda + (\mathfrak{B}_m \lambda) \lambda_l + (\mathfrak{B}_l \lambda) \lambda_m] C_{kh}^i \\ &+ \mathfrak{B}_l \mathfrak{B}_m [\vartheta(h_k^i C_h + h_{kh} C^i + h_h^i C_k)]. \end{aligned}$$

Using the condition (4.17) in above equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{kh}^i &= a_{lm} P_{kh}^i - a_{lm} \vartheta(h_k^i C_h + h_{kh} C^i + h_h^i C_k) + [\mathfrak{B}_l \mathfrak{B}_m \lambda + (\mathfrak{B}_m \lambda) \lambda_l \\ &+ (\mathfrak{B}_l \lambda) \lambda_m] C_{kh}^i + \mathfrak{B}_l \mathfrak{B}_m [\vartheta(h_k^i C_h + h_{kh} C^j + h_h^i C_k)]. \end{aligned}$$

Then Berwald's covariant derivative of the second order for the tensor  $\varphi(h_k^i C_h + h_{kh} C^i + h_h^i C_k)$  satisfies eq. (4.15) if and only if

$$\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i.$$

The above equation refer to  $P_{kh}^i$  satisfies the birecurrence property.

Taking  $\mathfrak{B}$ -covariant derivative for the condition (2.11) twice with respect to  $x^m$  and  $x^l$ , respectively, using eqs. (3.2) and (3.3) in the resulting equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jkh} &= \lambda a_{lm} C_{jkh} + [\mathfrak{B}_l \mathfrak{B}_m \lambda + (\mathfrak{B}_m \lambda) \lambda_l + (\mathfrak{B}_l \lambda) \lambda_m] C_{jkh} \\ &\quad + \mathfrak{B}_l \mathfrak{B}_m [\vartheta(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k)]. \end{aligned}$$

Using the condition (2.11) in above equation, we get

$$\begin{aligned} \mathfrak{B}_l \mathfrak{B}_m P_{jkh} &= a_{lm} P_{jkh} - a_{lm} \vartheta(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k) \\ &\quad + [\mathfrak{B}_l \mathfrak{B}_m \lambda + (\mathfrak{B}_m \lambda) \lambda_l + (\mathfrak{B}_l \lambda) \lambda_m] C_{jkh} \\ &\quad + \mathfrak{B}_l \mathfrak{B}_m [\vartheta(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k)]. \end{aligned}$$

Then Berwald's covariant derivative of the second order for the tensor  $\varphi(h_{jk} C_h + h_{kh} C_j + h_{hj} C_k)$  satisfies eq. (4.16) if and only if

$$\mathfrak{B}_l \mathfrak{B}_m P_{jkh} = a_{lm} P_{jkh}.$$

The above equation refer to  $P_{jkh}$  satisfies the birecurrence property. Hence, we have proved this theorem.

## 5. An Example

In this section, we give an example to clarify the proved findings.

**Example 5.1.** *The behavior of the torsion tensor  $P_{kh}^i$  as birecurrent if and only if the projection on indicatrix for it is also birecurrent.*

Firstly, since the torsion tensor  $P_{kh}^i$  behaves as birecurrent, then the condition (3.13) is satisfied. In view of (2.12), the projection of the torsion tensor  $P_{kh}^i$  on indicatrix is given by

$$p.P_{kh}^i = P_{bc}^a h_a^i h_k^b h_h^c. \quad (5.1)$$

Using  $\mathfrak{B}$ -covariant derivative for eq. (5.1) twice with respect to  $x^m$  and  $x^l$ , respectively, using the condition (3.13) and the fact that  $h_b^a$  is covariant constant in above equation, we get

$$\mathfrak{B}_l \mathfrak{B}_m (p.P_{kh}^i) = a_{lm} (P_{bc}^a h_a^i h_k^b h_h^c).$$

Using eq. (5.1) in above equation, we get

$$\mathfrak{B}_l \mathfrak{B}_m (p.P_{kh}^i) = a_{lm} (p.P_{kh}^i). \quad (5.2)$$

Equation (5.2) refers to the projection on indicatrix for the torsion tensor  $P_{kh}^i$  behaves as birecurrent.

Secondly, let the projection on indicatrix for the torsion tensor  $P_{kh}^i$  is birecurrent, i.e. satisfy eq. (5.2). Using (2.12) in eq. (5.2), we get

$$\mathfrak{B}_l \mathfrak{B}_m (P_{bc}^a h_a^i h_k^b h_h^c) = a_{lm} (P_{bc}^a h_a^i h_k^b h_h^c).$$

By using (2.13) in above equation, we get

$$\begin{aligned} & \mathfrak{B}_l \mathfrak{B}_m [P_{kh}^i - P_{kc}^i l^c l_h - P_{bh}^i l^b l_k + P_{bc}^i l^b l_k l^c l_h \\ & - P_{kh}^a l^i l_a + P_{kc}^a l^i l_a l^c l_h + P_{bh}^a l^i l_a l^b l_k - P_{bc}^a l^i l_a l^b l_k l^c l_h] \\ & = a_{lm} [P_{kh}^i - P_{kc}^i l^c l_h - P_{bh}^i l^b l_k + P_{bc}^i l^b l_k l^c l_h \\ & - P_{kh}^a l^i l_a + P_{kc}^a l^i l_a l^c l_h + P_{bh}^a l^i l_a l^b l_k - P_{bc}^a l^i l_a l^b l_k l^c l_h]. \end{aligned}$$

In view of (2.3) and if

$$P_{bc}^a y_a = P_{bc}^a y^b = P_{bc}^a y^c = 0,$$

then above equation can be written as

$$\mathfrak{B}_l \mathfrak{B}_m P_{kh}^i = a_{lm} P_{kh}^i.$$

The above equation means the torsion tensor  $P_{kh}^i$  behaves as birecurrent.

## 6. Conclusion

We obtained the necessary and sufficient condition for Cartan's second curvature tensor  $P_{jkh}^i$ , associate curvature tensor  $P_{ijkh}$ , torsion tensor  $P_{kh}^i$ ,  $P$ -Ricci tensor  $P_{jk}$ , curvature vector  $P_k$  and scalar curvature  $P$  which satisfy birecurrence property in  $\mathfrak{B}C - BRF_n$ ,  $P2$ -like  $-\mathfrak{B}C - BRF_n$ ,  $P^* - \mathfrak{B}C - BRF_n$  and  $P$ -reducible  $-\mathfrak{B}C - BRF_n$ . Furthermore, the relationship between Cartan's second curvature tensor  $P_{jkh}^i$  and  $(h)hv$ -torsion tensor  $C_{jk}^i$  in sense of Berwald has been discussed.

## REFERENCES

1. A. A. Abdallah, A. A. Navlekar and K. P. Ghadle, *On  $\mathfrak{B}$ -covariant derivative of first order for some tensors in different spaces*, J. Math. Analysis. Model. **2**(2) (2021), 30-37.
2. A. A. Abdallah, A. A. Navlekar and K. P. Ghadle, *Special types of generalized  $\mathfrak{B}P$ -recurrent spaces*, J. Computer. Math. Sci. **10**(5) (2019), 972-979.
3. A. A. Abdallah, A. A. Navlekar and K. P. Ghadle, *On  $P^*$ - and  $P$ -reducible of Cartan's second curvature tensor*, J. Appl. Sci. computation. **6**(11) (2019), 13-24.
4. A. A. Abdallah, A. A. Navlekar, K. P. Ghadle and A. A. Hamoud, *Decomposition for Cartan's second curvature tensor of different order in Finsler spaces*, Nonlinear. Functional. Analysis. Appl. **27**(2) (2022), 433-448.
5. A. A. Abdallah, A. A. Hamoud, A. A. Navlekar and K. P. Ghadle, *On special spaces of  $H(hv)$ -torsion tensor  $c_{jkh}$  in generalized recurrent finsler space*, Bull. Pure & Applied. Sci. Math. Stat. **41**(1) (2022), 74-80.
6. A. A. Saleem, *On certain generalized birecurrent and trirecurrent Finsler spaces*, M.Sc. Thesis, University of Aden, Yemen, 2011.

7. A. A. Saleem, and A. A. Abdallah, *Study On  $U^h$ - birecurrent Finsler spaces*, Int. J. Adv. Res. Sci. Commun. Tech. **2**(3) (2022), 28-39.
8. A. M. Hanballa, *On covariant differentiation of curvature tensors in Finsler spaces*, University of Aden, (Aden), (Yemen), 2016.
9. A. M. Otman, *On covariant differentiation for curvature tensor of third order in the sense of Berwald*, M.sc. Thesis, University of Aden, Yemen, 2018.
10. F. Y. Qasem and A. M. Hanballa, *On study  $K^h$ - generalized birecurrent Finsler space*, Int. J. Sci. Basic. Appl. Res. **25**(3), (2016), 208-216.
11. H. Izumi, *On  $P^*$ -Finsler space of scalar curvature*, Tensor N.S., **38**(1982), 220-222.
12. H. Rund, *The differential geometry of Finsler spaces*, Springer-Verlag, Berlin Göttingen, (1959); 2nd Edit. (in Russian), Nauka, (Moscow), 1981.
13. K. C. Sarangi and A. Goswami, *On  $C$ - recurrent Finsler space*, J. Int. Acad. Phys. Sci. **10** (2006), 17-24.
14. M. A. Alqufail, F. Y. Qasem and A. A. Muhib, *On study  $K^h$ - birecurrent Finsler spaces*, Sci. J. Fac. Edu. Thamar. 2015.
15. M. Dahl, *An brief introduction to Finsler geometry*, Springer, 2006.
16. M. Gheorghe, *The indicatrix in Finsler geometry*, Anal. Sti. Ale. Uuiv. Mat. Tomul LIII, (2007), 163-180.
17. M. Matsumoto, *On  $h$ -isotropic and  $C^h$ -recurrent Finsler*, J. Math. Kyoto Univ., **11** (1971), 1-9.
18. M. Matsumoto, *On Finsler spaces with curvature tensor of some special forms*, Tensor N.S., **22** (1971), 201-204.
19. P. K. Dwivedi,  *$P$ - reducible Finsler spaces and applications*, Int. J. Math. Analysis, **5**(5) (2011), 223-229.
20. P. N. Pandey and R. Verma,  *$C^h$ -birecurrent Finsler spaces*, J. Int. Acad. Phy. Sci. **2**(1998), 43-50.
21. P. N. Pandey and S. Dikshit, *On  $P^*$ - and  $P$ -reducible Finsler spaces of recurrent curvature*, J. Int. Acad. Phy. Sci. **2** (1998), 11-17.
22. P. S. Saxena and P. Swaroop, *A study of  $P^*$ -reducible Finsler space with Douglas tensor*, J. Int. Acad. Phy. Sci. **17**(3) (2013), 277-285.
23. S. Dikshit, *Certain types of recurrences in Finsler spaces*, D. phil. Thesis, University of Allahabad, (Allahabad), (India), 1992.
24. S. I. Ohta, *Comparison Finsler geometry*, Springer International Publishing, 2021.
25. S. M. Zamanzadeh, B. Najafi and M. Toomanian, *On generalized  $P$ -reducible Finsler manifolds*, Open. Math. **16**(2018), 718-723.

Received: 05.01.2023

Accepted: 22.05.2023